

部局	神戸大学大学院農学研究科
担当講座 (教育研究分野)	応用生命化学講座(食品・栄養化学教育研究分野)
就こうとする職名	教授
(ふりがな) 氏 名	(さかきばら ひろゆき) 榊原 啓之
性別	男
最終卒業学校名 学部学科等名 卒業(修了)年月	神戸大学 神戸大学大学院自然科学研究科生物機能化学専攻博士課程前期課程 修了 2002年3月31日
学位称号	博士(学術)(神戸大学) 2002年3月31日
学 歴	1997年3月31日 愛媛大学農学部生物資源学科 卒業 1997年4月1日 神戸大学大学院自然科学研究科生物機能化学専攻博士課程前期課程 入学 1999年3月31日 神戸大学大学院自然科学研究科生物機能化学専攻博士課程前期課程 修了 1999年4月1日 神戸大学大学院自然科学研究科生命科学専攻博士課程後期課程 入学 2002年3月31日 神戸大学大学院自然科学研究科生命科学専攻博士課程後期課程 修了 2002年3月31日 博士(学術)(神戸大学) 学位取得
職 歴	2022年2月27日 (株)小倉山本食品・非常勤研究員(～2002年5月) 2022年11月13日 神戸大学農学部特別研究員(～2003年1月) 2003年2月1日 フランス国立農学研究所(INRA/Clermont-Fd)博士研究員(～2004年2月) 2004年2月23日 神戸大学連携創造センター事務補佐員(～2004年3月) 2004年4月1日 神戸大学農学部特別研究員(～2005年1月) 2004年9月17日 武庫川女子大学非常勤講師(～2005年3月) 2005年2月1日 徳島大学大学院ヘルスバイオサイエンス研究部COE研究員(～2006年9月) 2006年10月1日 静岡県立大学助手(環境科学研究所)(～2007年3月31日) 2007年4月1日 静岡県立大学助教(環境科学研究所)(～2012年3月31日) 2009年1月10日 マックスプランク精神医学研究所(ドイツ・ミュンヘン)Visiting Researcher(～2009年3月) 2012年4月1日 宮崎大学准教授(農学部)(～2016年3月) 2016年4月1日 宮崎大学教授(農学部)

1. 寺尾純二, 下位香代子 (監修), 越阪部奈緒美, 榊原啓之, 中村宜督, 三好規之, 室田佳恵子 (編者), ポリフェノールの科学ー基礎化学から健康機能までー, (株)朝倉書店、東京 2023 年 11 月.
2. 宇田津徹朗, 河原聡, 榊原啓之 (著者), 平田昌彦 (編者), 生物・農学系のための統計学-大学での基礎学修から研究論文まで- (第 2 版), 主担当: 2 つの正規変量間の関係, 朝倉書店, 217 ページ, 2023 年.
3. 榊原啓之, 下位香代子, 抗疲労・抗ストレス・睡眠改善食品の開発 (監修: 井上和生, 山崎英恵), 抗ストレス機能の評価法, シーエムシー出版, 287 ページ, 担当 69~77 ページ, 2020 年.
4. 中村宜督, 榊原啓之, 室田佳恵子 (著者/編者), エッセンシャル食品化学, 主担当: ビタミンとミネラル (ミネラル), 視覚成分 (天然色素), 嗅覚成分 (天然香気成分), 講談社, 247 ページ, 主担当 92~102, 129~136, 167~174 ページの執筆および全体校正, 2018 年.
5. 榊原啓之, Visual 栄養学テキスト 食べ物と健康 I 食品学総論 食品の成分と機能 (監修: 寺尾純二, 村上明), 香気・におい成分, 中山書店, 148 ページ, 担当 90-95 ページ, 2018 年.
6. 榊原啓之, 下位香代子, 機能性食品開発のための初期評価試験プロトコール集 (監修: 山本 (前田) 万里), 単独隔離ストレス負荷試験 (動物試験), 培養細胞を用いた抗ストレス作用評価試験 I (受容体競合試験), 培養細胞を用いた抗ストレス作用評価試験 II (シグナル伝達・遺伝子発現・浸潤試験), シーエムシー出版, 366 ページ, 担当 223~231 ページ, 2017 年.
7. 宇田津徹朗, 河原聡, 榊原啓之 (著者), 平田昌彦 (編者), 生物・農学系のための統計学-大学での基礎学修から研究論文まで-, 主担当: 2 つの正規変量間の関係, 朝倉書店, 217 ページ, 主担当: 91~107 ページおよび全体校正, 2017 年.
8. 榊原啓之, 下位香代子, 寺尾純二, 植物色素フラボノイド (監修: 武田幸作, 齊藤規夫, 岩科司), フラボノイドの薬理作用, 文一総合出版, 695 ページ, 担当 590~600 ページ, 2013 年.
9. 榊原啓之, 下位香代子, 寺尾純二, 抗ストレス食品の開発と展望 II (監修: 横越英彦), 抗ストレス食品の動物による機能評価, シーエムシー出版, 295 ページ, 担当 29~40 ページ, 2012 年.
10. Ohura T. and Sakakibara H., Polycyclic Aromatic Hydrocarbons: Pollution, Health Effects and Chemistry (Polymer Science and Technology) (編: Haines PA and Hendrickson MD), Chlorinated polycyclic aromatic hydrocarbons: A new class of environmental toxin, Nova Biomedical, 361 ページ, 担当 107~112 ページ, 2009 年.
11. 下位香代子, 榊原啓之, 食品に関して知っておくべき衛生安全知識のすべて『食品安全ハンドブック』(編: 食品安全ハンドブック編集委員会編), 各種変異原性からみた食品関連の有機化学物質 (第 8 章 有害化学物質による汚染-8.1), 丸善出版, 864 ページ, 担当 590~600 ページ, 2010 年.
12. Sakakibara H., Ohura T., Morita M., Hirabayashi S., Kuruto-Niwa R., Amagai T., and Shimoi K.,

- Interdisciplinary Studies on Environmental Chemistry-Environmental Research in Asia for Establishing a Scientist's Network (Eds: Obayashi Y. et al.), Aryl hydrocarbon receptor-mediated gene expression by chlorinated polycyclic aromatic hydrocarbons and cross-talk with estrogen receptors, TERRAPUB Publishing, 341 ページ, 担当 197~202 ページ, 2009 年.
13. Morita M., Ohura T., Kuruto-Niwa R., Sakakibara H., Amagai T., and Shimoi K., Persistent Organic Pollutants (POPs) Research in Asia (Ed: Morita, M.), Chlorinated phenanthrenes Induce CYP1A1 through aryl hydrocarbon receptor activation, DIOXIN2007 Organizing Committee, 583 ページ, 担当 302~307 ページ, 2008 年.
14. Sakakibara H., Izawa Y., Nakajima J., Seo S., Tamaki T., Kawai Y., and Terao J., ACS symposium series 993 Functional Food and Health (Eds: Shibamoto T et al.), Effects of polyphenol rich herbal medicine, Ginkgo biloba extracts on neurotransmitter levels in rat brain, Oxford University Press, 514 ページ, 担当 429~434 ページ, 2008 年.
15. 寺尾純二, 榊原啓之; 抗ストレス食品の開発と展望 (監修: 横越英彦), 人参サポニン, 羅布麻エキス, シーエムシー出版, 338 ページ, 担当 222~226 ページ, 238~242 ページ, 2006 年.

1. Ismayawati A.D.<sup>#</sup>, Yokoyama D., Nishiguchi Y.<sup>#</sup>, Saeki Y., Sakakibara H.<sup>\*</sup>, Effect of harvest time of day on the isoflavone synthesis pathway in immature soybeans, *Food Research*, in press.
2. Suzuki, Y.<sup>#</sup>, Yokoyama, D., Matsuura, C., Kondo, K., Shimazaki, T., Ryoike, K., Kobayashi, A. and Sakakibara, H.<sup>\*</sup>, Active-phase plasma alkaline phosphatase isozyme activity is a sensitive biomarker for excessive fructose intake. *In Vivo*, 2023, 107, 105652.
3. Kubota, H.<sup>#</sup>, Shinohara, S.<sup>#</sup>, Yokoyama, D., Tanaka, W.<sup>#</sup>, Matsuyama, H.<sup>#</sup>, Arimura, T., Kimura, S. and Sakakibara, H.<sup>\*</sup>, Suppressive effects of sugarcane molasses concentrate on starch-induced hyperglycemia in mice. *Journal of Functional Foods*, **107**, 105652, 2023. (計 8 ページ)
4. Tanaka, W.<sup>#</sup>, Matsuyama, H.<sup>#</sup>, Shimoi, K., Yokoyama, D. and Sakakibara, H.<sup>\*</sup>, Social confrontation stress decreases hepatic fibroblast growth factor-21 expression in aged mice. *Biochemistry and Biophysics Reports*, **34**, 101454, 2023.
5. Suzuki, Y.<sup>#</sup>, Kondo, K., Toyoda, K., Tanaka, Y., Kobayashi, A., Yokoyama, D. and Sakakibara, H.<sup>\*</sup>, Novel biomarker establishment for evaluation of excessive fructose consumption using a rat model. *In Vivo*, **37 (1)**, 173–181, 2023.
6. Tajiri, H.<sup>#</sup>, Tanaka, W.<sup>#</sup>, Matsuyama, H.<sup>#</sup>, Sugita, T., Hidaka, K., Yokoyama, D. and Sakakibara, H.<sup>\*</sup>, Beneficial effects of the consumption of hot-water extracts of thinned immature mangos (*Mangifera indica* "Irwin") on the hypertriglyceridemia of apolipoprotein E-deficient mice. *Metabolites*, **12 (2)**, 116, 2022.
7. Shibazaki, N.<sup>#</sup>, Tanaka, W.<sup>#</sup>, Suzuki, S.<sup>#</sup>, Matsuyama, H.<sup>#</sup>, Kubota, H.<sup>#</sup>, Ogawa, K., Yokoyama, D. and Sakakibara, H.<sup>\*</sup>, Maltose consumption exacerbates high-fat diet-induced overweight and related parameters in mice. *Functional Foods in Health and Disease*, **12 (11)**, 680–691, 2022.
8. Mizuno, M., Fujioka, A., Bitani, S., Minato, K.I. and Sakakibara, H., Anti-allergic activity of fucoidan can be enhanced by coexistence with quercetin. *International Journal of Molecular Sciences*, **23 (20)**, 12163, 2022.
9. Matsuzaka, H.<sup>#</sup>, Matsuyama, H.<sup>#</sup>, Tanaka, W.<sup>#</sup>, Tajiri, H.<sup>#</sup> and Sakakibara, H.<sup>\*</sup>, Selective consumption of fish oil at end of the day increases the physiological fatty acid compositions of eicosapentaenoic acid and docosahexaenoic acid in mice. *Molecules*, **27 (4)**, 1271, 2022.
10. Takashima, M.<sup>#</sup>, Tanaka, W.<sup>#</sup>, Matsuyama, H.<sup>#</sup>, Tajiri, H.<sup>#</sup> and Sakakibara, H.<sup>\*</sup>, Maternal quercetin consumption during pregnancy may help regulate total cholesterol/HDL-cholesterol ratio without effect on cholesterol levels in male progeny consuming high-fat diet. *Nutrients*, **13 (4)**, 1242, 2021.
11. Tajiri, H.<sup>#</sup>, Tanaka, W.<sup>#</sup>, Takashima, M.<sup>#</sup>, Matsuyama, H.<sup>#</sup>, Sugita, T., Hidaka, K. and

- Sakakibara, H.<sup>\*</sup>, Subchronic safety evaluation of hot-water extract from thinned immature mangos (*Mangifera indica* 'Irwin'): 90-days oral toxicity study in rats. *Toxicology Reports*, **8**, 1046–1053, 2021.
12. Matsuyama, H.<sup>#</sup>, Tanaka, W.<sup>#</sup>, Miyoshi, N., Miyazaki, T., Michimoto, H. and Sakakibara, H.<sup>\*</sup>, Beneficial effects of the consumption of sun-dried radishes (*Raphanus sativus* cv. YR-Hyuga-Risou) on dyslipidemia in apolipoprotein E-deficient mice. *Journal of Food Biochemistry*, **45** (5), e13727, 2021.
13. Arimura, T., Watanabe, S., Sakakibara, H. and Kimura, S., The new sugar to suppress postprandial hyperglycemia: Improvement of refined white sugar by mixing specially grown brown cane extract. *Acta Scientific NUTRITIONAL HEALTH*, **5.6**, 43–50, 2021.
14. Yasuda, M.T., Watanabe, Y., Hosono, T., Seki, T., Suzuki, T.A., Sakakibara, H. and Shimoi, K., Nocturnal light exposure stimulates the cardiac fibrinolysis system and stress responses in C3H/He mice. *Thrombosis Research*, **188**, 79–81, 2020.
15. Tsuboi, H., Sakakibara, H., Matsunaga, M., Tatsumi, A., Yamakawa-Kobayashi, K., Yoshida, N. and Shimoi, K., Omega-3 eicosapentaenoic acid Is related to happiness and a sense of fulfillment-A study among female nursing workers. *Nutrients*, **12** (11), 3462, 2020.
16. Tanaka, W.<sup>#</sup>, Matsuyama, H.<sup>#</sup>, Yokoyama, D., Yamashita, Y., Ashida, H., Sakono, S. and Sakakibara, H.<sup>\*</sup>, Daily consumption of black soybean (*Glycine max* L.) seed coat polyphenols attenuates dyslipidemia in apolipoprotein E-deficient mice. *Journal of Functional Foods*, **72**, 104054, 2020.
17. Nakanishi, T., Kagamizono, K., Yokoyama, S., Suzuki, R., Sakakibara, H., Sugamoto, K., Erickson, L. and Kawahara, S., Dietary phytanic acid-induced changes in tissue fatty acid profiles in mice. *Journal of Dairy Research*, **87** (4), 498–500, 2020.
18. Nakanishi, T., Kagamizono, K., Yokoyama, S., Suzuki, R., Sakakibara, H., Erickson, L. and Kawahara, S., Effects of dietary phytol on tissue accumulation of phytanic acid and pristanic acid and on the tissue lipid profiles in mice. *Animal Science Journal*, **91** (1), e13424, 2020.
19. Nagase, S., Karashima, S., Tsujiguchi, H., Tsuboi, H., Miyagi, S., Kometani, M., Aono, D., Higashitani, T., Demura, M., Sakakibara, H., Yoshida, A., Hara, A., Nakamura, H., Takeda, Y., Nambo, H., Yoneda, T. and Okamoto, S., Impact of gut microbiome on hypertensive patients with low-salt intake: Shika study results. *Frontiers in Medicine*, **7**, 475, 2020.
20. Matsuyama, H.<sup>#</sup>, Tanaka, W.<sup>#</sup>, Yokoyama, D., Matsumoto, S., Sano, T., Yamashita, T., Nishimura, S., Sakono, M. and Sakakibara, H.<sup>\*</sup>, Suitability of a 10% fat diet for use in time-restricted feeding experiments with C57BL/6 mice. *Bioactive Compounds in Health*

- and Disease*, **3**, 55–65, 2020.
21. Friedman, M., Sakakibara, H., Mizuno, M., Kim, D.H. and Kozukue, N., Free amino acid, phenolic, flavonoid,  $\beta$ -carotene, lycopene, dehydrotomatine, and  $\alpha$ -tomatine content of peel powders prepared from commercial cherry tomatoes. *Current Topics in Phytochemistry*, **16**, 1–16, 2020.
  22. Tsuboi, H., Sakakibara, H., Tatsumi, A., Yamakawa-Kobayashi, K., Matsunaga, M., Kaneko, H. and Shimoi, K., Serum IL-6 levels and oxidation rate of LDL cholesterol were related to depressive symptoms independent of omega-3 fatty acids among female hospital and nursing home workers in Japan. *Journal of Affective Disorders*, **249**, 385–393, 2019.
  23. Tanaka, W.<sup>#</sup>, Yokoyama, D., Matsuura, Y.<sup>#</sup>, Nozaki, M., Hirozawa, N., Kunitake, H., Sakono, M. and Sakakibara, H.<sup>\*</sup>, Subchronic toxicity evaluation of leaves from rabbiteye blueberry (*Vaccinium virgatum* Aiton) in rats. *Toxicology Reports*, **6**, 272–278, 2019.
  24. Itoh, K., Masumori, S., Mukai, D., Sakakibara, H., Yasuda, M. and Shimoi, K., Dosage time affects alkylating agents induced micronuclei in mouse peripheral blood reticulocytes through the function of erythropoietin. *The Journal of Toxicological Sciences*, **44** (4), 273–282, 2019.
  25. Friedman, M., Kozukue, N., Mizuno, M., Sakakibara, H., Choi, S.H., Fujitake, M. and Land, K.M., The analysis of the content of biologically active phenolic compounds, flavonoids, and glycoalkaloids in harvested red, yellow, and green tomatoes, tomato leaves, and tomato stems. *Current Topics in Phytochemistry*, **15**, 43–53, 2019.
  26. Yokoyama, D.<sup>#</sup>, Tanaka, W.<sup>#</sup>, Hashizume, Y., Tandia, M., Sakono, M., Shimoi, K. and Sakakibara, H.<sup>\*</sup>, Daily consumption of monoglucosyl rutin prevents high-fat diet-induced obesity by suppressing gastric inhibitory polypeptide secretion in mice. *Functional Foods in Health and Disease*, **8** (7), 353–371, 2018.
  27. Tsuboi, H., Sakakibara, H., Minamida, Y., Tsujiguchi, H., Matsunaga, M., Hara, A. and Nakamura, H., Elevated levels of serum IL-17A in community-dwelling women with higher depressive symptoms. *Behavioral Sciences*, **8** (11), 102, 2018.
  28. Nohara, C.<sup>#</sup>, Yokoyama, D., Tanaka, W.<sup>#</sup>, Sogon, T., Sakono, M. and Sakakibara, H.<sup>\*</sup>, Daily Consumption of Bilberry (*Vaccinium myrtillus* L.) Extracts Increases the Absorption Rate of Anthocyanins in Rats. *Journal of Agricultural and Food Chemistry*, **66** (30), 7958–7964, 2018.
  29. Matsuura, Y.<sup>#</sup>, Sakakibara, H., Kawaguchi, M., Murayama, E., Yokoyama, D., Yukizaki, C., Kunitake, H. and Sakono, M., Effects of blueberry leaf and stem extracts on hepatic lipid levels in rats consuming a high-sucrose diet. *Functional Foods in Health and Disease*, **8** (9), 447–461, 2018.

30. Tashiro, K.<sup>#</sup>, Sakakibara, H., Yokoyama, D.<sup>#</sup>, Uehara, T., Matsuura, Y. and Sakono, M., Consumption of salted pickles of sun-dried radish roots (*Raphanus sativus* cv. YR-Hyuga-Risou) attenuates blood pressure in spontaneously hypertensive rats. *Food Science and Technology Research*, **23 (5)**, 757–763, 2017.
31. Horii, Y., Nagasawa, T., Sakakibara, H., Takahashi, A., Tanave, A., Matsumoto, Y., Nagayama, H., Yoshimi, K., Yasuda, M.T., Shimoi, K. and Koide, T., Hierarchy in the home cage affects behaviour and gene expression in group-housed C57BL/6 male mice. *Scientific Reports*, **7 (1)**, 6991, 2017.
32. Yokoyama, D.<sup>#</sup>, Sakakibara, H., Fukuno, H., KKimura, K., Hori, A., Nara, T., Kato, K. and Sakono, M., Whey protein-hydrolyzed peptides diminish hepatic lipid levels on rats consuming high-sucrose diets. *Food Science and Technology Research*, **22**, 631–638, 2016.
33. Miura, A., Sugiyama, C., Sakakibara, H., Shimoi, K. and Goda, T., Bioavailability of isoflavones from soy products in equol producers and non-producers in Japanese women. *Journal of Nutrition and Intermediary Metabolism*, **6**, 41–47, 2016.
34. Ohura, T., Sakakibara, H., Watanabe, I., Shim, W.J., Manage, P.M. and Guruge, K.S., Spatial and vertical distributions of sedimentary halogenated polycyclic aromatic hydrocarbons in moderately polluted areas of Asia. *Environmental Pollution*, **196**, 331–340, 2015.
35. Yamazaki, S., Sakakibara, H., Takemura, H., Yasuda, M. and Shimoi, K., Quercetin-3-O-glucuronide inhibits noradrenaline binding to alpha2-adrenergic receptor, thus suppressing DNA damage induced by treatment with 4-hydroxyestradiol and noradrenaline in MCF-10A cells. *The Journal of Steroid Biochemistry and Molecular Biology*, **143**, 122–129, 2014.
36. Sakakibara, H.<sup>\*</sup>, Ohura, T., Kamiya, Y., Yamanaka, N., Shimada, N., Shimoi, K. and Guruge, K.S., Sex-dependent difference in the hepatic and pulmonary toxicological effects in mice administrated 7-chlorinated benz[*a*]anthracene. *Fundamental Toxicological Sciences*, **1 (3)**, 101–108, 2014.
37. Sakakibara, H., Ichikawa, Y., Tajima, S., Makino, Y., Wakasugi, Y., Shimoi, K., Kobayashi, S., Kumazawa, S. and Goda, T., Practical application of flavonoid-poor menu meals to the study of the bioavailability of bilberry anthocyanins in human subjects. *Bioscience, Biotechnology, and Biochemistry*, **78 (10)**, 1748–1752, 2014.
38. Kato, A., Sakakibara, H., Tsuboi, H., Tatsumi, A., Akimoto, M., Shimoi, K., Ishii, T., Kaneko, H., Nakayama, T. and Ohashi, N., Depressive symptoms of female nursing staff working in stressful environments and their association with serum creatine kinase and lactate dehydrogenase - a preliminary study. *BioPsychoSocial Medicine*, **8**, 21, 2014.

39. Katayama, M., Donai, K., Sakakibara, H., Ohtomo, Y., Miyagawa, M., Kuroda, K., Kodama, H., Suzuki, K., Kasai, N., Nishimori, K., Uchida, T., Watanabe, K., Aso, H., Isogai, E., Sone, H. and Fukuda, T., Coffee consumption delays the hepatitis and suppresses the inflammation related gene expression in the Long-Evans Cinnamon rat. *Clinical Nutrition*, **33 (2)**, 302–310, 2014.
40. Aoshima, Y.<sup>#</sup>, Sakakibara, H., Suzuki, T.A., Yamazaki, S. and Shimoi, K., Nocturnal light exposure alters hepatic *Pai-1* expression by stimulating the adrenal pathway in C3H mice. *Experimental Animals*, **63 (3)**, 331–338, 2014.
41. Shimamura, Y., Yoda, M., Sakakibara, H., Matsunaga, K. and Masuda, S., Pu-erh tea suppresses diet-induced body fat accumulation in C57BL/6J mice by down-regulating SREBP-1c and related molecules. *Bioscience, Biotechnology, and Biochemistry*, **77 (7)**, 1455–1460, 2013.
42. Sakakibara, H., Ohura, T., Kido, T.<sup>#</sup>, Yamanaka, N., Tanimura, N., Shimoi, K. and Guruge, K.S., Organ-specific distribution of 7-chlorinated benz[*a*]anthracene and regulation of selected cytochrome P450 genes in rats. *The Journal of Toxicological Sciences*, **38 (1)**, 137–143, 2013.
43. Kido, T.<sup>#</sup>, Sakakibara, H., Ohura, T., Guruge, K.S., Kojima, M., Hasegawa, J., Iwamura, T., Yamanaka, N., Masuda, S., Sakaguchi, M., Amagai, T. and Shimoi, K., Evaluation of chlorinated benz[*a*]anthracene on hepatic toxicity in rats and mutagenic activity in *Salmonella typhimurium*. *Environmental Toxicology*, **28 (1)**, 21–30, 2013.
44. 榊原啓之, 市川陽子, 多島早奈英, 牧野尚恵, 若杉悠佑, 熊澤茂則, 佐々木敏, 下位香代子 and 合田敏尚, 日本人の食事摂取基準 (2005年版) に基づいたフラボノイド低減食の作成方法の提案. *日本栄養・食糧学会誌*, **37 (5)**, 229–235, 2012.
45. Yamazaki, S., Sakakibara, H., Takemura, H. and Shimoi, K., 4-Hydroxyestradiol induces  $\gamma$ -H2AX in the presence of an inhibitor of catechol-O-methyltransferase in human breast cancer MCF-7 cells. *Genes and Environment*, **34 (3)**, 129–135, 2012.
46. Sakakibara, H., Suzuki, A.<sup>#</sup>, Kobayashi, A., Motoyama, K., Matsui, A., Sayama, K., Kato, A., Ohashi, N., Akimoto, M., Nakayama, T. and Shimoi, K., Social isolation stress induces hepatic hypertrophy in C57BL/6J mice. *The Journal of Toxicological Sciences*, **37 (5)**, 1071–1076, 2012.
47. Okada, Y., Yi, Y., Suzuki, T., Sakuragawa, S., Sakakibara, H., Shimoi, K. and Sugiura, T., Wearable ECG recorder with acceleration sensors for monitoring daily stress. *Journal of Medical and Biological Engineering*, **34**, 129–135, 2012.



48. Itoh, K., Masumori, S., Nakajima, M., Hayashi, M., Sakakibara, H. and Shimoi, K., Differences in micronucleus induction in peripheral blood reticulocytes of mice exposed to N-ethyl-N-nitrosourea at light and dark dosing times. *The Journal of Toxicological Sciences*, **37 (2)**, 427–430, 2012.
49. Yoshino, S., Hara, A., Sakakibara, H., Kawabata, K., Tokumura, A., Ishisaka, A., Kawai, Y. and Terao, J., Effect of quercetin and glucuronide metabolites on the monoamine oxidase-A reaction in mouse brain mitochondria. *Nutrition*, **27 (7-8)**, 847–852, 2011.
50. Tsuboi, H., Sakakibara, H., Yamakawa-Kobayashi, K., Tatsumi, A., Inamori, T., Hamamoto, R., Suzuki, A. and Shimoi, K., Val1483Ile polymorphism in the fatty acid synthase gene was associated with depressive symptoms under the influence of psychological stress. *Journal of Affective Disorders*, **134 (1-3)**, 448–452, 2011.
51. Sakakibara, H., Romanowski, C.P., Jakubcakova, V., Flachskamm, C., Shimoi, K. and Kimura, M., Feeble awake effects of plasminogen activator inhibitor type-1 in mice. *Behavioural Brain Research*, **220 (2)**, 354–357, 2011.
52. Kobayashi, A., Yokoyama, H., Kataoka, J., Ishida, T., Kuno, H., Sugai, S., Sakakibara, H. and Shimoi, K., Effects of spaced feeding on gene expression of hepatic transaminase and gluconeogenic enzymes in rats. *The Journal of Toxicological Sciences*, **36 (3)**, 325–337, 2011.
53. Ishisaka, A., Ichikawa, S., Sakakibara, H., Piskula, M.K., Nakamura, T., Kato, Y., Ito, M., Miyamoto, K., Tsuji, A., Kawai, Y. and Terao, J., Accumulation of orally administered quercetin in brain tissue and its antioxidative effects in rats. *Free Radical Biology and Medicine*, **51 (7)**, 1329–1336, 2011.
54. Agawa, S.<sup>#,§</sup>, Sakakibara, H.<sup>§</sup>, Iwata, R., Shimoi, K., Hergesheiber, A. and Kumazawa, S., Anthocyanins in mesocarp/epicarp and endocarp of fresh açai (*Euterpe oleracea* Mart) and their antioxidant activity and bioavailability. *Food Science and Technology Research*, **17 (4)**, 327–334, 2011.
- §Equally contribution
55. Takemura, H., Uchiyama, H., Ohura, T., Sakakibara, H., Kuruto, R., Amagai, T. and Shimoi, K., A methoxyflavonoid, chrysoeriol, selectively inhibits the formation of a carcinogenic estrogen metabolite in MCF-7 breast cancer cells. *The Journal of Steroid Biochemistry and Molecular Biology*, **118 (1-2)**, 70–76, 2010.
56. Takemura, H., Nagayoshi, H., Matsuda, T., Sakakibara, H., Morita, M., Matsui, A., Ohura, T. and Shimoi, K., Inhibitory effects of chrysoeriol on DNA adduct formation with benzo[*a*]pyrene in MCF-7 breast cancer cells. *Toxicology*, **274 (1-3)**, 42–48, 2010.

57. Takemura, H., Itoh, T., Yamamoto, K., Sakakibara, H. and Shimoi, K., Selective inhibition of methoxyflavonoids on human CYP1B1 activity. *Bioorganic & Medicinal Chemistry*, **18 (17)**, 6310–6315, 2010.
58. Sakakibara, H., Koyanagi, A.<sup>#</sup>, Suzuki, T.A., Suzuki, A., Ling, L. and Shimoi, K., Effects of animal care procedures on plasma corticosterone levels in group-housed mice during the nocturnal active phase. *Experimental Animals*, **59 (5)**, 637–642, 2010.
59. Ohura, T., Morita, M., Kuruto-Niwa, R., Amagai, T., Sakakibara, H. and Shimoi, K., Differential action of chlorinated polycyclic aromatic hydrocarbons on aryl hydrocarbon receptor-mediated signaling in breast cancer cells. *Environmental Toxicology*, **25 (2)**, 180–187, 2010.
60. Kobayashi, A., Oshida, S., Yamazaki, Y., Maekawa, T., Kuno, H., Sugai, S., Sakakibara, H. and Shimoi, K., Relationships between plasma and tissue transaminase activities in rats maintained under different feeding conditions. *The Journal of Toxicological Sciences*, **35 (5)**, 639–652, 2010.
61. Kato, A., Naitou, H., Namioka, M., Akimoto, M., Ishii, T., Sakakibara, H., Shimoi, K., Nakayama, T. and Ohashi, N., Proteomic identification of serum proteins associated with stress-induced gastric ulcers in fasted rats. *Bioscience, Biotechnology, and Biochemistry*, **74 (4)**, 812–818, 2010.
62. Kageyama, A.<sup>#</sup>, Sakakibara, H., Zhou, W., Yoshioka, M., Ohsumi, M., Shimoi, K. and Yokogoshi, H., Genistein regulated serotonergic activity in the hippocampus of ovariectomized rats under forced swimming stress. *Bioscience, Biotechnology, and Biochemistry*, **74 (10)**, 2005–2010, 2010.
63. Yokoyama, A.<sup>§</sup>, Sakakibara, H.<sup>§</sup>, Crozier, A., Kawai, Y., Matsui, A., Terao, J., Kumazawa, S. and Shimoi, K., Quercetin metabolites and protection against peroxynitrite-induced oxidative hepatic injury in rats. *Free Radical Research*, **43 (10)**, 913–921, 2009.

<sup>§</sup>Equally contribution

64. Sakakibara, H., Ogawa, T., Koyanagi, A., Kobayashi, S., Goda, T., Kumazawa, S., Kobayashi, H. and Shimoi, K., Distribution and excretion of bilberry anthocyanins in mice. *Journal of Agricultural and Food Chemistry*, **57 (17)**, 7681–7686, 2009.
65. Sakakibara, H., Miyashita, T.<sup>#</sup>, Matsui, A., Ozaki, A., Suzuki, A., Ikeda, T., Morishita, K. and Shimoi, K., Effects of St John's Wort extracts on corticosterone levels dysregulated by social isolation stress. *Journal of Traditional Medicines*, **26 (2)**, 86–92, 2009.
66. Motoyama, K., Nakai, Y., Miyashita, T., Fukui, Y., Morita, M., Sanmiya, K., Sakakibara,

- H., Matsumoto, I., Abe, K., Yakabe, T., Yajima, N. and Shimoi, K., Isolation stress for 30 days alters hepatic gene expression profiles, especially with reference to lipid metabolism in mice. *Physiological Genomics*, **37** (2), 79–87, 2009.
67. Kobayashi, A., Suzuki, Y., Kuno, H., Sugai, S., Sakakibara, H. and Shimoi, K., Effects of fenofibrate on plasma and hepatic transaminase activities and hepatic transaminase gene expression in rats. *The Journal of Toxicological Sciences*, **34** (4), 377–387, 2009.
68. Sakakibara, H., Yoshino, S., Miyazaki, T., Kawai, Y. and Terao, J., Antidepressant-like effect of Ginseng (*Panax ginseng* C. A. Meyer) in behavioral models. *Journal of Clinical Biochemistry and Nutrition*, **43** (Suppl 1), 255–258, 2008.
69. Sakakibara, H., Yoshino, S., Kawai, Y. and Terao, J., Antidepressant-like effect of onion (*Allium cepa* L.) powder in a rat behavioral model of depression. *Bioscience, Biotechnology, and Biochemistry*, **72** (1), 94–100, 2008.
70. Ogawa, K.<sup>#,§</sup>, Sakakibara, H.<sup>§</sup>, Iwata, R., Ishii, T., Sato, T., Goda, T., Shimoi, K. and Kumazawa, S., Anthocyanin composition and antioxidant activity of the Crowberry (*Empetrum nigrum*) and other berries. *Journal of Agricultural and Food Chemistry*, **56** (12), 4457–4462, 2008.
- <sup>§</sup>Equally contribution
71. Hashimoto, T., Goto, M., Sakakibara, H., Oi, N., Okamoto, M. and Kanazawa, K., Yellow tea is more potent than other types of tea in suppressing liver toxicity induced by carbon tetrachloride in rats. *Phytotherapy Research*, **21** (7), 668–670, 2007.
72. Sakakibara, H., Ishida, K., Grundmann, O., Nakajima, J., Seo, S., Butterweck, V., Minami, Y., Saito, S., Kawai, Y., Nakaya, Y. and Terao, J., Antidepressant effect of extracts from Ginkgo biloba leaves in behavioral models. *Biological and Pharmaceutical Bulletin*, **29** (8), 1767–1770, 2006.
73. Sakakibara, H., Ashida, H., Fukuda, I., Furuyashiki, T., Sano, T., Nonaka, Y., Hashimoto, T. and Kanazawa, K., A frequent drinking of green tea lowers the levels of endogenous oxidative stress in small intestines, erythrocytes and kidneys in rats. *Journal of Clinical Biochemistry and Nutrition*, **39** (1), 32–39, 2006.
74. Hashimoto, T., Ueda, Y., Oi, N., Sakakibara, H., Piao, C., Ashida, H., Goto, M. and Kanazawa, K., Effects of combined administration of quercetin, rutin, and extract of white radish sprout rich in kaempferol glycosides on the metabolism in rats. *Bioscience, Biotechnology, and Biochemistry*, **70** (1), 279–281, 2006.
75. Hashimoto, T., Oi, N., Sakakibara, H., Goto, M. and Kanazawa, K., Comparative study of antioxidative activity among the six great chinese teas; green, white, yellow, oolong,

- black, and pu-erh teas. *ITE Letters on Batteries, New Technologies & Medicine*, **7 (5)**, 484–487, 2006.
76. Sakakibara, H., Nakagawa, S., Wakameda, H., Nakagiri, Y., Kamata, K., Das, S.K., Tsuji, T. and Kanazawa, K., Effects of Japanese kelp (kombu) on life span of benzo[*a*]pyrene-fed mice. *Journal of Nutritional Science and Vitaminology*, **51 (5)**, 369–373, 2005.
77. Sakakibara, H., Ishida, K., Izawa, Y., Minami, Y., Saito, S., Kawai, Y., Butterweck, V., Tamaki, T., Nakaya, Y. and Terao, J., Effects of forced swimming stress on rat brain function. *The Journal of Medical Investigation*, **52 Suppl**, 300–301, 2005.
78. Sakakibara, H., Viala, D., Ollier, A., Combeau, A. and Besle, J.M., Isoflavones in several clover species and in milk from goats fed clovers. *Biofactors*, **22 (1-4)**, 237–239, 2004.
79. Ashida, H., Furuyashiki, T., Nagayasu, H., Bessho, H., Sakakibara, H., Hashimoto, T. and Kanazawa, K., Anti-obesity actions of green tea: possible involvements in modulation of the glucose uptake system and suppression of the adipogenesis-related transcription factors. *Biofactors*, **22 (1-4)**, 135–140, 2004.
80. Sakakibara, H., Honda, Y., Nakagawa, S., Ashida, H. and Kanazawa, K., Simultaneous determination of all polyphenols in vegetables, fruits, and teas. *Journal of Agricultural and Food Chemistry*, **51 (3)**, 571–581, 2003.
81. Sakakibara, H., Ashida, H. and Kanazawa, K., A novel method using 8-hydroperoxy-2'-deoxyguanosine formation for evaluating antioxidative potency. *Free Radical Research*, **36 (3)**, 307–316, 2002.
82. Sun, M., Sakakibara, H., Ashida, H., Danno, G. and Kanazawa, K., Dietary antioxidants fail in protection against oxidative genetic damage in in vitro evaluation. *Bioscience, Biotechnology, and Biochemistry*, **64 (11)**, 2395–2401, 2000.
83. Sun, M., Sakakibara, H., Ashida, H., Danno, G. and Kanazawa, K., Cytochrome P4501A1-inhibitory action of antimutagenic anthraquinones in medicinal plants and the structure-activity relationship. *Bioscience, Biotechnology, and Biochemistry*, **64 (7)**, 1373–1378, 2000.
84. Saeki, K., Sakakibara, H., Sakai, H., Kunito, T. and Tanabe, S., Arsenic accumulation in three species of sea turtles. *Biometals*, **13 (3)**, 241–250, 2000.
85. Kanazawa, K. and Sakakibara, H., High content of dopamine, a strong antioxidant, in Cavendish banana. *Journal of Agricultural and Food Chemistry*, **48 (3)**, 844–848, 2000.

1. Yamashita, Y., Sakakibara, H., Toda, T. and Ashida, H., Insights into the potential benefits of black soybean (*Glycine max* L.) polyphenols in lifestyle diseases. *Food and Function*, **11 (9)**, 7321–7339, 2020.
2. Sakakibara, H. and Shimoi, K., Anti-stress effects of polyphenols: animal models and human trials. *Food and Function*, **11 (7)**, 5702–5717, 2020.
3. 榊原啓之, 下位香代子, 社会的ストレスに対するフラボノイド含有食品の効果. *日本ポリフェノール学術雑誌*, **6 (1)**, 11–14, 2017.
4. Torii-Yasuda, M., Sakakibara, H. and Shimoi, K., Estrogen- and stress-induced DNA damage in breast cancer and chemoprevention with dietary flavonoid. *Genes and Environment*, **39**, 10, 2017.
5. Sakakibara, H., Torii-Yasuda, M. and Shimoi, K., Effects of environmental and social stressors on biological rhythms. *The Journal of Physical Fitness and Sports Medicine*, **5 (2)**, 143–152, 2016.
6. 榊原啓之, 青島良輝#, 山崎隼輔, 下位香代子, 栄養素の消化・吸収機構に関与している体内時計. *New Food Industry*, **55 (12)**, 1–8, 2013.
7. Tsurusaki, T.#, Sakakibara, H.\*, Aoshima, Y., Yamazaki, S., Sakono, M. and Shimoi, K., Diurnal rhythmicity in biological processes involved in bioavailability of functional food factors. *Journal of Clinical Biochemistry and Nutrition*, **52 (3)**, 208–214, 2013.
8. Takemura, H., Sakakibara, H., Yamazaki, S. and Shimoi, K., Breast cancer and flavonoids - a role in prevention. *Current Pharmaceutical Design*, **19 (34)**, 6125–6132, 2013.
9. 榊原啓之, 小柳顯陽#, 青島良輝#, 鈴木敬明, 木村昌由美, 下位香代子, 外部刺激に対する生体応答を評価するために一夜行性のマウスを用いた時間生物学的アプローチ. *谷本学校毒性質問箱*, **第13号**, 116–123, 2011. (査読なし)
10. 榊原啓之, 寺尾純二, 抗うつ様活性を有する食品成分および漢方薬の検索. *FFI Journal of Japan*, **211**, 721–726, 2006.